

Inflationary cosmology - a dissipative quantum field theory process

Arjun Berera*

School of Physics, University of Edinburgh EH9 3JZ, Great Britain

To appear in Proceedings ICHEP02, Amsterdam, July 2002

The current status of early universe cosmology can be summarized as precision, but inherently limited, data requiring explanation from precision and robust theory. Specifically inflation is presently the single concrete hope for solving the early universe problem, since it is generically realizable in quantum field theory, but treatment of interactions needs considerable attention. This talk reviews the importance and progress in this direction.

I. INTRODUCTION

Early universe cosmology is a subject with limited access to observation, since there is only so much information that can be acquired about this time period from our one local region in the universe. As such, a convincing theory about this time period must rely substantially on the soundness and unambiguity of its mathematical foundation. At present, inflation is the most hopeful idea for developing into a theory, since it is based on quantum field theory, and quantum fields so far have been exclusively successful in explaining the high energy world, up to at least the TeV scale. However, consistent solutions of inflation founded on quantum field theory have been allusive. Inflationary dynamics inherently is a multifield problem, since the vacuum energy that drives inflation eventually must convert to radiation, which generally is comprised of a variety of particle species. In the earliest conception of inflation [1], it was pictured that inflation would result in a isentropic expansion that would rapidly put the universe in a supercooled thermodynamic phase. Subsequently it was observed that supercooled inflation is not mandatory and that nonisentropic inflationary expansion, warm inflation, also is possible [2]. Moreover, since the main condition for inflation is $\rho_v > \rho_r$, where ρ_v, ρ_r are the vacuum and radiation energy densities respectively, supercooled inflation appears as a limiting case within the general regime of warm inflation.

The most nontrivial aspect of the inflaton model is understanding the energy transfer dynamics from potential energy to radiation. In the most commonly followed inflation picture, dissipative effects of the inflaton field are

ignored throughout the inflation period, leading to a supercooled inflationary regime. However, from a thermodynamic perspective, this picture appears very restrictive. The point being, even if the inflaton were to allow a minuscule fraction of the energy to be released, say one part in 10^{20} , it still would constitute a significant radiation energy density component in the universe. For example, for inflation with vacuum (i.e. potential) energy at the GUT scale $\sim 10^{15-16}$ GeV, leaking one part in 10^{20} of this energy density into radiation corresponds to a temperature of 10^{11} GeV, which is nonnegligible. In fact, the most relevant lower bound that cosmology places on the temperature after inflation comes from the success of hot Big-Bang nucleosynthesis, which requires the universe to be within the radiation dominated regime by $T \gtrsim 1$ GeV. This limit can be met in the above example by dissipating as little as one part in 10^{60} of the vacuum energy density into radiation. Thus, from the perspective of both interacting field theory and basic notions of equipartition, it appears to be a highly tuned requirement of supercooled inflation to prohibit the inflaton from such tiny amounts of dissipation.

These considerations have led to examining the possibility of warm inflation, an inflationary regime in which radiation also is present. Warm inflation is comprised of non-isentropic expansion in the background cosmology [2] and thermal seeds of density perturbations [3,4] (for related earlier work please see [5]). During warm inflation, interactions between the inflaton and other fields cause the radiation energy density to remain substantial due to its constant production from conversion of vacuum energy. This expansion regime is intrinsically different from the supercooled inflation regime, since warm inflation smoothly terminates into a subsequent radiation dominated regime, without a reheating period.

The warm inflation picture has one immediate conceptual advantage, that the dynamics is free of questions about quantum-to-classical transition. The scalar inflaton field is in a classical state, thus justifying the application of a classical evolution equation, and the inflaton fluctuations, which induce the metric perturbations, are classical. Another problem alleviated by dissipative effects is the initial condition problem of inflation [6].

Warm inflation continues proving to be a promising solution to the early universe problem, with now several variants and perspectives on the basic idea [4,7]. Development of it has had two directions, which will be addressed in this talk. First is deriving a first principle quantum field theory realization of its dynamics. Second

*PPFAC Advanced Fellow

is determining observational tests for warm inflation.

II. FIRST PRINCIPLES ORIGIN

The earliest works looked for high temperature warm inflation solutions, under rigid adiabatic, equilibrium conditions [8]. Within this limited framework, one type of warm inflation solution was obtained [8]. The high-T regime was examined first, since considerable methodology was already available for treating it. However, intrinsically, the statistical state relevant for warm inflation is not required to be an equilibrium state. The slowly varying nature of the macroscopic variables in warm inflation cosmology suggest that the statistical state may not be far from equilibrium, although this is something that should be proven from the dynamics. Much work remains in order to develop the mathematical formalism necessary to address this problem. As one step in this direction to fill the missing gaps, recently we studied the zero temperature dissipative dynamics of interacting scalar field systems in Minkowski spacetime [9] (for another interesting direction see [10]). This is useful to understand, since the zero temperature limit constitutes a baseline effect that will be prevalent in any general statistical state. The key result presented in this talk is that for a broad range of cases, involving interaction with as few as one or two fields, we find dissipative regimes for the scalar field system. This is important for inflationary cosmology, since it suggests that dissipation may be the norm not exception for an interacting scalar field system, thus warm inflation could be a natural dynamics once proper treatment of interactions is done.

We study the Lagrangian, where

$$\mathcal{L}_I = -\frac{\lambda}{4!}\Phi^4 - \sum_{j=1}^{N_\chi} \left\{ \frac{f_j}{4!}\chi_j^4 + \frac{g_j^2}{2}\Phi^2\chi_j^2 \right\} - \sum_{k=1}^{N_\psi} \bar{\psi}_k \left[h_{k\phi}\Phi + \sum_{j=1}^{N_\chi} h_{kj\chi}\chi_j \right] \psi_k \quad (1)$$

and $\Phi \equiv \varphi + \phi$ such that $\langle \Phi \rangle = \varphi$. Our aim is to obtain the effective equation of motion for $\varphi(t)$ and from that determine the energy dissipated from the $\varphi(t)$ system into radiation.

The tadpole method, which requires $\langle \phi \rangle = 0$, gives effective equation of motion for $\varphi(t)$,

$$\ddot{\varphi}(t) + m_\phi^2\varphi(t) + \frac{\lambda}{6}\varphi^3(t) + \frac{\lambda}{2}\varphi(t)\langle\phi^2\rangle + \frac{\lambda}{6}\langle\phi^3\rangle + \sum_{j=1}^{N_\chi} g_j^2 [\varphi(t)\langle\chi_j^2\rangle + \langle\phi\chi_j^2\rangle] + \sum_{k=1}^{N_\psi} h_{k\phi}\langle\bar{\psi}_k\psi_k\rangle = 0. \quad (2)$$

The field expectation values in this equation are obtained by solving the coupled set of field equations. In our cal-

culational, we have evaluated them in a perturbative expansion using dressed Green's functions. One general feature of these expectation values is they will depend of the causal history of $\varphi(t)$, so that Eq. (2) is a temporally nonlocal equation of motion for $\varphi(t)$.

The general expression for the effective equation of motion is given in [9] and is very complicated. Formally, we can examine Eq. (2) within a Markovian-adiabatic approximation, in which the equation of motion is local in time and the motion of $\varphi(t)$ is slow. At $T = 0$, such an approximation is not rigorously valid. Nevertheless, this approximation allows understanding the magnitude of dissipative effects. Furthermore, we have shown in [9] that the nonlocal effects tend to filter only increasingly higher frequency components of $\varphi(t)$ from nonlocal effects increasingly further back in time. Thus for low frequency components of $\varphi(t)$, memory only is retained to some short interval in the past. Since within the adiabatic approximation, $\varphi(t)$ only has low frequency components, we believe the Markovian-adiabatic approximation is legitimate at least for order of magnitude estimates. Within this approximation, the effective equation of motion for $\varphi(t)$ has the general form

$$\ddot{\varphi}(t) + m_\phi^2\varphi(t) + \frac{\lambda}{6}\varphi^3(t) + \eta(\varphi)\dot{\varphi}(t) = 0, \quad (3)$$

where

$$\eta(\varphi) \approx \varphi^2(t) \frac{N_\psi}{256\pi^2} \left(\frac{\lambda^2 h_{\phi\psi}^2}{4m_\phi} + \frac{N_\chi g^4 h_{\chi\psi}^2}{m_\chi} \right).$$

An alternative to the above Lagrangian based derivation, is the canonical derivation following formalism developed in the mid 80's [11]. Although the canonical and Lagrangian approaches should agree, the former is far less developed in dissipative quantum field theory, in particular for treating interactions. Nevertheless, the canonical approach provides useful insight, especially for understanding the origin of particle creation. For example, consider in the canonical approach the expectation value of $\langle\chi_i^2\rangle$,

$$\int \frac{d^3q}{(2\pi)^3 2\text{Re}\omega_{\mathbf{q},\chi_j}} [2x_{\mathbf{q},\chi_j}(t) + 2\text{Re}y_{\mathbf{q},\chi_j}(t) + 1],$$

where $x_{\mathbf{q},\chi_j}(t) = \langle a_{\mathbf{q},\chi_j}^\dagger(t)a_{\mathbf{q},\chi_j}(t) \rangle$ is the particle number density and $y_{\mathbf{q},\chi_j}(t) = \langle a_{\mathbf{q},\chi_j}(t)a_{-\mathbf{q},\chi_j}(t) \rangle$ is the off-diagonal correlation. The evolution equations for $x_{\mathbf{q},\chi_j}(t)$ and $y_{\mathbf{q},\chi_j}(t)$ can be obtained from the field equation for χ_j to give

$$\dot{x}_{\mathbf{q},\chi_j} = \frac{\dot{\omega}_{\mathbf{q},\chi_j}}{\omega_{\mathbf{q},\chi_j}} \text{Re} y_{\mathbf{q},\chi_j}, \quad \dot{y}_{\mathbf{q},\chi_j} = \frac{\dot{\omega}_{\mathbf{q},\chi_j}}{\omega_{\mathbf{q},\chi_j}} \left[x_{\mathbf{q},\chi_j} + \frac{1}{2} \right] - 2i\omega_{\mathbf{q},\chi_j} y_{\mathbf{q},\chi_j}. \quad (4)$$

To yield dissipation, it is noted in [11] that the correlation amongst produced particles needs to be destroyed sufficiently rapidly and $\dot{x}_{\mathbf{q},\chi_j}(t)$ and $\langle\chi_j(t)\rangle$ should become local functions of time involving $\varphi(t)$ and $\dot{\varphi}(t)$. Based on this requirement, [11] asserts that $\omega_{\mathbf{q},\chi_j}$ in the above equation for $\dot{y}_{\mathbf{q},\chi_j}(t)$ must have an imaginary part, which in fact should be the χ_j -particle decay width, $\text{Im}\omega_{\mathbf{q},\chi_j} \propto \Gamma_{\mathbf{q},\chi_j}$. Applying these assumptions, the contribution to $\eta(\varphi)$ in Eq. (3) from the χ_j field, say $\eta_{\chi_j}(\varphi)$, once again is obtained (up to $\mathcal{O}(1)$ factors). Since this approach has ad-hoc assumptions, it still is incomplete and requires development. Nevertheless, the approach is interesting and for now accepting the assumptions, the origin of particle creation and energy conservation are clearly seen. In particular, the particle production rate is given by $\int (d^3k/(2\pi)^3) \dot{x}_{\mathbf{q},\chi_j}(t) \omega_{\mathbf{q},\chi_j}$ and similar to [11], it can be shown this is equal to the vacuum energy loss rate from the χ_j field contribution, $\eta_{\chi_j}(\varphi)\dot{\varphi}^2$.

Returning to Eq. (3), estimated magnitudes of energy production will be obtained in the overdamped regime of warm inflation, $m^2(\phi) = m_\phi^2 + \lambda\phi^2/2 < \eta^2$. In this regime, the energy dissipated by the scalar field goes into radiation energy density ρ_r at the rate $\dot{\rho}_r = -dE_\phi/dt = \eta(\varphi)\dot{\varphi}^2$.

In [9] we have determined radiation production for two cases (a). $m(\varphi) > m_\chi > 2m_\psi$, (b). $m_\chi > 2m_\psi > m(\varphi)$. To focus on a case typical for inflation, suppose the potential energy is at the GUT scale $V(\varphi)^{1/4} \sim 10^{15}\text{GeV}$ and we consider the other parameters in a regime consistent with the e-fold and density fluctuation requirements of inflation. Note, although this is a flat nonexpanding spacetime analysis, since the dissipative effects will be at subhorizon scale, one expects these estimates to give a reasonable idea of what to expect from a similar calculation done in expanding spacetime. Expressing the radiation in terms of a temperature scale as $T \sim \rho_r^{1/4}$, we find for case (a) $1\text{GeV} < T < 10^7\text{GeV} < H$ and for case (b) $T \gtrsim 10^{14}\text{GeV} > H$, where $H = \sqrt{8\pi V/(3m_p^2)}$.

III. OBSERVATIONAL TESTS

Supercooled inflation has three parameters, related to the potential energy magnitude V_0 , slope ϵ , and curvature η , whereas there are four observable constraints (δ_H , A_g , n_s , n_g). This implies a redundancy in the observations and allows for a consistency relation [12]. This is usually expressed as a relationship between the tensor-to-scalar ratio and the slope of the tensor spectrum. Warm inflation has an extra parameter, the dissipation factor, which implies four constraints for four parameters. Hence we do not expect the consistency relation of standard inflation to hold in warm inflation [13]. Thus, to discriminate between warm and standard inflation, it requires a measure of all four observables. The upcoming MAP

and Planck satellite missions should provide strong constraints on the scalar spectrum and being equipped with polarization detectors, it is hoped the tensor spectrum also will be measured. Recently nongaussian effects from warm inflation models were computed and found to be too small to measure [14].

IV. CONCLUSION

The key point emphasized in this talk has been that any typical inflaton field theory model has interactions and so dissipation can be an unexceptional consequence. These dissipative effects have always been ignored, unjustifiably, in supercooled inflation, but as shown here their effects can be nontrivial. The most significant effect of dissipation is to completely alter the inflationary regime from supercooled to warm. However, irrespective of the inflation regime, as discussed above and in [6], dissipative effects before the inflation period may be important in alleviating the initial condition problem. Also, dissipative effects imply new realizations of thermal inflation [15]. Moreover, dissipative effects can alter density perturbations. In this talk we discussed the case where dissipation leads to an ideal thermalized state, thus with thermal density perturbations. However the opposite limit can be considered from ideal thermalization to that of negligible influence from the radiation field on the density perturbations, so that only the quantum fluctuations contribute. Then simple calculations show warm inflation models would *not* require fine tuning, so that scalar potentials of reasonably large curvature would be acceptable. The nature of density perturbations during warm inflation still requires much investigation in particular to understand the interplay between radiation and the scalar field. In conclusion, this talk discussed how dissipation during inflation could be robust and possibly solve inflation's most challenging dilemma, the scalar field fine tuning problem.

-
- [1] A. H. Guth, Phys. Rev. **D23** (1981) 347; K. Sato, Phys. Lett. B **99** (1981) 66; A. Linde, Phys. Lett. B **108** (1982) 389; A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48** (1982) 1220.
 - [2] A. Berera, Phys. Rev. Lett. **75** (1995) 3218; Phys. Rev. D **54** (1996) 2519; Phys. Rev. D **55** (1997) 3346.
 - [3] A. Berera and L. Z. Fang, Phys. Rev. Lett. **74** (1995) 1912; A. Berera, Nucl. Phys. B **585** (2002) 666.
 - [4] H. P. De Oliveira and S. E. Joras, Phys. Rev. D **64** (2001) 063513; H. P. De Oliveira, Phys. Lett. B **526** (2002) 1; L. P. Chimento *et. al.*, Phys. Rev. D **65** (2002) 083510.

- [5] I. Moss, Phys. Lett. B **154** (1985) 120; R. Brout and S. P. Spindel, Nucl. Phys. B **348** (1991) 405; S. P. Spindel and R. Brout, Phys. Lett. B **320** (1994) 241.
- [6] A. Berera and C. Gordon, Phys. Rev. D **63** (2001) 063505; R. O. Ramos, Phys. Rev. D **64** (2001) 123510.
- [7] H. P. de Oliveira and R. O. Ramos, Phys. Rev. **D57** (1998) 741; J. M. F. Maia and J. A. S. Lima, Phys. Rev. **D60** (1999) 101301; W. Lee and L.-Z. Fang, Phys. Rev. **D59** (1999) 083503.
- [8] A. Berera, M. Gleiser and R. O. Ramos, Phys. Rev. D **58** (1998) 123508; Phys. Rev. Lett. **83** (1999) 264.
- [9] A. Berera and R. O. Ramos, Phys. Rev. D **63** (2001) 103509.
- [10] I. D. Lawrie, Phys. Rev. D **66** (2002) 041702.
- [11] M. Morikawa and S. Sasaki, Prog. Theor. Phys. **72** (1984) 782.
- [12] A. R. Liddle and D. H. Lyth, Phys. Rept. **231** (1993) 1.
- [13] A. N. Taylor and A. Berera, Phys. Rev. D **62** (2000) 083517.
- [14] S. Gupta *et. al.*, Phys. Rev. **D66** (2002) 043510.
- [15] A. Berera and T. W. Kephart, hep-ph/0207208.